

# PREDICTORS AND CONSEQUENCES OF SUBSTANCE USE AMONG PREGNANT CIGARETTE SMOKERS

Research Assistants: Nicolas Thor and Millie Pal; Principle Investigator: Dr. Xiaozhong Wen MD, PhD  
Division of Behavioral Medicine, Department of Pediatrics, UB Jacobs School of Medicine and Biomedical Sciences

## Objectives

- To investigate:**
- Predictors for substance use among pregnant cigarette smokers (i.e., co-use of tobacco and substance)
  - The effects of co-use on success in smoking cessation
  - The effects of co-use on maternal and infant health

## Background

- Co-use of tobacco and marijuana is common among teens and young adults (Ramo 2012)
- Marijuana is the most commonly used (illicit) substance during pregnancy (Metz 2015)
- As the main psychoactive compound of marijuana, tetrahydrocannabinol (THC) is able to cross the placenta as well as contaminate breastmilk in expectant and lactating mothers, respectively (Jaques 2014)
- Marijuana use throughout pregnancy is associated with growth restriction in mid- and late pregnancy, leading to low birth weight (Jaques 2014)

## Methods

- **Sample:** 56 pregnant smokers from UB Pregnancy and Smoking Cessation Study (2015-2017; Buffalo, NY)
- **Measurements:** substance use, socio-demographics, breastfeeding intention (self-report via survey), smoking status (urine cotinine), anthropometrics (scales)
- **Data analysis:** ANOVA and Chi-square tests in SAS 9.3 software

### Study flowchart




Figure 1. Study flowchart

## Results

### Substance Use Among Pregnant Smokers




Figure 2. Substance use among pregnant smokers

- Marijuana was the most commonly used substance during pregnancy (47.5%), followed by cocaine (3.1%).

Table 1. Socio-demographic, pregnancy, and smoking characteristics of the eligible sample (N=56)

Characteristic	n (%)	Mean (SD)
Age, years		27.9 (5.6)
≤ 24	19 (33.9)	
25-29	16 (28.6)	
≥ 30	21 (37.5)	
Race/ethnicity		
Non-Hispanic, Caucasian	14 (25.0%)	
Non-Hispanic, African American	33 (58.9%)	
Hispanic or American Indian	9 (16.1%)	
Education level		
High school or lower	28 (50.0%)	
Some college or vocational training	20 (35.7%)	
2-year or 4 year college degree	8 (14.3%)	
Married	16 (28.6%)	
Employed	19 (33.9%)	
Household income, US dollars		
< 5,000	19 (33.9%)	
5,000-11,999	13 (23.2%)	
12,000-24,999	12 (21.4%)	
≥ 25,000	12 (21.4%)	
Gestation at enrollment, weeks		
≤ 13	31 (55.4%)	
24-27	23 (41.1%)	
≥ 28	2 (3.6%)	
Number of cigarettes smoked, per day		
1-4	11 (19.6%)	
5-9	24 (42.9%)	
≥ 10	21 (37.5%)	

Figure 3. Sample Characteristics

### Marijuana Co-Use




Figure 4. Proportion of marijuana co-user among pregnant cigarette smokers

- 77% of pregnant cigarette smokers in our study have used marijuana at least once in their lives
- 47.5% continued to use marijuana during their pregnancy

### Change in Marijuana Use Throughout the Smoking Cessation Intervention




Figure 5. Change in marijuana use throughout the smoking cessation intervention

- After our intervention on cigarette smoking cessation, only 12.5% still used marijuana
- Further decreased to 6.3% by the end of pregnancy

### Predictors for Co-Use of Marijuana Among Pregnant Smokers




Figure 6. Predictors for co-use of marijuana among pregnant smokers. P ≤ 0.05

- 60.0% of light smokers (1-4 cigs/day) also used marijuana during pregnancy, which was much higher compared to moderate (5-9 cigs/day; 27.6%) and heavy smokers (10+ cigs/day; 31.3%)
- 73.7% with some college education used marijuana during pregnancy, which was much higher than those with completion of ≤ high school (25.0%) or a college degree (44.4%)
- 100% (vs 28.6%) of the participants who believed that their current partner has a drug problem used marijuana during pregnancy

### Consequences of Co-use of Marijuana on Gestational Weight Gain




Figure 7. Consequences of co-use of marijuana on gestational weight gain

- Although not significant, marijuana users had higher mean gestational weight gain than non-users

### Breastfeeding Intention




Figure 8. Breastfeeding intention

- Also not significant, but interesting to point out, cessation of marijuana use before pregnancy was associated with higher breastfeeding intention

## Conclusion

- Majority of our participants have reported marijuana use in their lifetime. However, most of them quit marijuana as our cigarette smoking cessation intervention and their pregnancies progressed.
- Light cigarette smoking, some college education, and having partner with drug problems predict co-use of marijuana during pregnancy among cigarette smokers.
- The co-use of marijuana and cigarettes did not predict gestational weight gain and breastfeeding intention.
- The co-use of marijuana and cigarettes did not predict other maternal and infant outcomes including the mother's success in quitting smoking cigarettes and sleep quality; infant's gestational age, weight, length, or body mass index at birth. (data not shown)

## References

1. Agrawal, A., Budney, A. J., & Lynskey, M. T. (2012). The co-occurring use and misuse of cannabis and tobacco: a review. *Addiction, 107*(7), 1221-1233.
2. Day, N. L., Goldschmidt, L., & Thomas, C. A. (2006). Prenatal marijuana exposure contributes to the prediction of marijuana use at age 14. *Addiction, 101*(9), 1313-1322.
3. Jaques, S. C., et al. (2014). Cannabis, the pregnant woman and her child: weeding out the myths. *Journal of Perinatology, 34*(6), 417-42
4. Metz, T. D., & Stickrath, E. H. (2015). Marijuana use in pregnancy and lactation: a review of the evidence. *American journal of obstetrics and gynecology, 213*(6), 761-778.
5. Miller, D. S., & Miller, T. Q. (1997). A test of socioeconomic status as a predictor of initial marijuana use. *Addictive Behaviors, 22*(4), 479-489.
6. Ramo, D. E., Liu, H., & Prochaska, J. J. (2012). Tobacco and marijuana use among adolescents and young adults: a systematic review of their co-use. *Clinical psychology review, 32*(2), 105-121.

## Acknowledgement

- **Collaborators:** Dr. Leonard H. Epstein, Dr. Stephen T. Higgins, Dr. Rina D. Eiden, Dr. Faye E. Justicia-Linde, Dr. Youfa Wang, Dr. Kai Ling Kong, Rocco A. Paluch
- **Research Assistants**
- **Health professionals at recruitment sites**
- **All patients and their families**

## Funding

- NIH CTSA Pilot Fund, UB Dept of Pediatrics, CURCA

## Contact

Dr. Xiaozhong Wen, MD, PhD; Assistant Professor (716-829-6811; xiaozhon@buffalo.edu)